

# Vomizal®

## Antiemetic

### Composition :

- Each tablet of VOMIZAL® contains:  
Meclozine hydrochloride 25 mg  
Pyridoxine hydrochloride 50 mg

### Indications:

- Nausea and vomiting of pregnancy .
- Treatment of emesis induced by motion sickness, other drugs and in anesthesia.
- Treatment of vertigo associated with diseases affecting the vestibular system .

### Dosage and administration:

- For severe nausea and vomiting of pregnancy :  
One VOMIZAL® tablet in the morning and one tablet at bed time.
- For motion sickness: take one tablet one hour before traveling ,to be repeated every 24 hours as needed .
- For post operative vomiting: Two tablets four hours before surgery .
- Anti vertigo agent : one to four tablets as needed in divided doses .

### Contraindications:

- VOMIZAL® is contraindicated in:  
- Patients on levodopa therapy .
- A Previous hypersensitivity to any components of this product .

### Side Effects:

- Blurred vision .
- Drowsiness .
- Dryness of mouth , nose and throat .

### Precautions:

- Because of drowsiness and possible sedative effect of VOMIZAL® so drive should be avoided .
- VOMIZAL® should be given with caution to patients with impaired liver function .
- VOMIZAL® should be given with caution to patients with asthma , glaucoma , peptic ulcer and prostatic hypertrophy.

### Drug Interactions:

- VOMIZAL® enhances the sedative effects of CNS depressants including barbiturates , hypnotics, opioid analgesics , anxiolytic

sedatives ,anti psychotics and alcohol .

- VOMIZAL® may suppress the cutaneous histamine response to allergen extracts and should be stopped for at least 72 hours before skin testing .

### Pregnancy and Lactation:

- In pregnancy : VOMIZAL® does not cause adverse effects on pregnant woman or her foetus ,so advice to use it in nausea and vomiting with pregnancy .
- In lactation :  
Meclozine hydrochloride may be excreted in breast milk and may inhibits lactation due to its anti muscarinic action .

### Properties:

- VOMIZAL® is anti emetic and anti nauseant .It diminishes vestibular stimulation and depresses labyrinthine function , also acts on medullary chemoreceptor trigger zone.
- Meclozine hydrochloride has a sedating anti histaminic with anti muscarinic and moderate sedative properties so it is used in the prevention and treatment of nausea and vomiting associated with a variety of conditions.
- Pyridoxine has benefits in pregnancy due to it's conversion in the erythrocytes to pyridoxal phosphate -the active form of the vitamine -which acts as coenzyme for numerous metabolic functions affecting protein , carbohydrates and lipid utilization .

### Packaging :

Blister of 10 tablets , pack of 2 blisters.

### Storage:

- Store in dry and cool place, at temperature below 30° C.

#### THIS IS A MEDICATION

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of pharmacist who sold the medication .
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consultation your doctor.
- Keep medication out of reach of children.

Produced by:  
Azal Pharmaceutical Industries Co.Ltd.  
Sudan - Khartoum



الأبعاد: 17×11 cm

نوع الخط: Times New Roman

حجم الخط: ٧.٥

رقم البانتون: 7427 C

تاريخ التعديل: ٢٠١٣/١/١٤ هـ

	Date	Sign
R&D		
Q.C		
Q.A.		
Tec. Manager		